

THE FIRST GLANCE

ABDOMINOPLASTY AND LIPOSUCTION POST CARE INSTRUCTIONS

Supplies to have at home:

- Your prescription medications (will be given to you on surgery day if not at pre-op appt).
- Gravol (oral medication & suppositories available). Used to treat nausea.
- Stool softener (ie: Restorolax/Lax-a-Day/Senokot). To prevent constipation. *Start day after surgery
- Antibiotic Ointment - (Original)
- Flexible gauze wrap (ie: Johnson & Johnson 3" or 4" soft gauze roll) Non-stick gauze or nursing pads work well.
- Hypoallergenic Paper Medical Tape (3M or similar)
- Kotex or similar sanitary napkins (pads) or gauze
- **IF FLANK LIPO-** Depends (or similar adult diaper), bring day of surgery to wear home (drainage from lipo sites is normal) and change as needed for first few days
- Band-aids
- Garbage bag and old towels to protect the seat of the car on your ride home
- Gatorade/Powerade or similar.
- Abdominal binder/medical compression garment- sized day of Pre-op.

*If garment is given to you day of pre-op, Pls bring it back with you on day of surgery.

Rest and Relaxation:

- When resting, it is ok to lie on your side. Tuck your legs up when lying on your side and place 2 to 3 pillows under your knees when sleeping on your back.

***IF MINI TUMMY TUCK-** you can sleep with only 1 pillow under knees when sleeping on your back

- Take 10 nice big deep breaths every hour while awake.
- When coughing or sneezing open your mouth and place your hands on your abdomen. Gently push on your abdominal incision to protect it.

Medications:

- Take your pain medication as directed when you get home. You may find these do not eliminate your discomfort 100%, but you will feel more comfortable. Pain is expected, the goal is to make the pain tolerable.
- Take all medications as prescribed, ensure you finish your entire course of antibiotics even if you do not experience any signs of infection.
- Resume prescribed medications unless otherwise stated.
- Follow all instructions which may accompany the medications such as: taking with food, avoiding alcohol and driving restrictions.

THE FIRST GLANCE

Activities:

- Make sure you walk as much as possible and lean forward at your waist (approx 30° angle) to decrease tension on your abdominal incision.
- After 2 weeks you can begin to stand straight.
*IF MINI TUMMY TUCK- you do not have to walk leaning forward.
- If you have had your rectus (abdominal) muscles tightened, do not lift more than 10 lbs or perform abdominal exercises for minimum of 4 weeks.
- Wait at least 2 weeks before participating in physical activities that will increase your heart rate and blood pressure.

Washing

- You may shower once the outer dressings are removed (even if you still have drains in.) It may be helpful to wear a lanyard around your neck and pin the drains up to keep them out of the way.
- Dry surgical site by gently patting with a clean towel or allow to air dry.
- Do not soak surgical sites - no baths, pools, hot tubs, or lake water for 4 weeks.
- You do not require any bandages once your initial bandages are removed, however, most patients feel more comfortable with a bandage to provide extra protection. Flexible gauze wrap can be used as a dressing once the original dressings have been removed.

Dressings/Wound Care

- Outer White Dressing may be removed after 24hrs.
- Apply Antibiotic ointment to Umbilicus
- Leave your steri-strips (paper tapes on incisions) on for as long as possible, typically they can be removed around 2 weeks. If the edges start to lift, you can trim them or if they are very saturated or are starting to come off sooner then you may remove them.
- Once the steri-strips are removed, apply Polysporin to your incisions once a day for one week.
- It is normal to experience some redness and irritation to the incision around your navel during the first week or two, this is due to the type of suture that is used in the area.
- Bleeding from your surgical site may occur following surgery. If it occurs hold constant firm pressure for 10 minutes with a clean towel.

THE FIRST GLANCE

- Dressings to Liposuction sites can also be removed at 24hrs post-op. Drainage of excess fluid is normal after lipo for several days. Have Kotex pad/Incontinent pads/Diapers or Thick gaze at home to change dressings as needed. Wear clothing that you do not mind if they get soiled.
- You may be sent home with Foam. It can be applied to the liposuction areas once drainage has stopped as directed by Dr. Islur.

Garments

- If you DID require abdominal muscle repair (sheath plication) Wear your abdominal binder/garment for 20 out of 24 hours per day for the first 4 weeks), then for 12 hours during the day for 2 more weeks.
- If you DID NOT require muscle repair then wear your garment for 20hrs per day for 2 weeks then for 12 hrs during the day for 2 more weeks
- The garment will help to aid in skin retraction, control swelling and reduce your risk of complications.
- When removing the garment, do so lying down as releasing the compression could make some people light headed.

****Binder should reach a few inches below the incision line****

- Take time to rest and recover from your surgery, have a healthy diet and get lots of rest. It is important that you follow these recommendations carefully so we can attain your surgical goals

Lymphatic Massage of the areas is recommended, this may start 3-4 days post-op, or as soon as leakage/drainage from incisions stop. Plan to make massage appts prior to surgery,

Drain Care

- Empty drains as instructed and document drainage. Drains are removed once drainage is less than 30cc over 24 hours per side. Abdominal drains must stay in for a minimum of 3 days.
- If you are from out-of-town (4+ hours away) your drains will be removed once drainage is less than 30cc over a 24-hour period PLUS 2 days, this is to further reduce the incidence of seroma requiring drainage.

Anticipate your drains being in for approximately 10 days, but no longer than 14 days.

- Apply Polysporin to drain site and cover with gauze to absorb fluid that may leak out. This is normal and can occur for 1 week after drain have been removed
- Once removed, you can shower 12 hrs after drain removal. Drain sites will close in 7 to 10 days.
- If there is a gush of fluid that occurs, this is likely fluid that was not picked up by the drain, this is normal.
- Get into the shower and gently push on your abdomen or breast to encourage the fluid to drain out.
- If drainage persists or is predominantly fresh blood, contact Dr. Islur.

1851 Grant Avenue, Winnipeg, MB, R3N 1Z2

(204) 885-1569 | www.thefirstglance.ca | contactus@thefirstglance.ca

THE FIRST GLANCE AESTHETIC CLINIC & SURGERY CENTRE

THE FIRST GLANCE

For all Liposuction Procedures

- Bruising tends to last about 2 weeks. Swelling can be significant and initial subsiding can take up to 6-8 weeks and up to 10 months for the final result to show. Swelling can increase during menstrual cycle.
- Recovery varies in every individual. Most feel uncomfortable for the first couple of days and depending on the occupation and extent of the surgery, can return to work within 1 week post-op

Three Weeks after your Surgery

- Begin to massage your incisions with Scar Recovery Gel/Stick
- If your incisions are going to be exposed to the sun, apply SPF 45 or higher for at least one year to prevent tanning.

*Do not sleep with pets in the bed to avoid infection

*No baths, hot tubs, swimming pools, and lake water for 3 weeks or until your incisions are fully healed in order to avoid infection

*Avoid exposure to second-hand smoke for at least the first few weeks after surgery

THE FIRST GLANCE

Please contact the clinic immediately and request to speak to a nurse or physician if you experience any of the following after surgery.

- Increased pain, increased redness, creamy or pus-like discharge, foul odour, or
- Increased warmth around surgical site, rash, fever exceeding 38.8° C (101.8°F)
- Chest pain (for continuous chest pain lasting more than 5 minutes – call 911)
- Unexplained sudden onset of shortness of breath
- Rapid pulse or heart palpitations
- Severe pain in leg that does not resolve and/or swelling of leg
- Changes in skin colour of an extremity, such as turning pale, red or blue
- Dizziness that does not resolve, or fainting
- Coughing up blood / bloody diarrhea

After Hour Care

If there is a concern about your surgery or your healing, please call The First Glance Clinic at 204-885-1419.

If the clinic is closed (evenings or weekends) or in case of an emergency, please call Dr. Islur - 431-302-5177

If after 3 attempts you cannot reach Dr Islur, please call the plastic surgery resident by calling St Boniface Hospital Operator at 204-237-2053

Visit www.drislur.ca for more info